

514 Rec'd PCT/PTO 13 MAR 2000

FORM PTO 1390 (REV 5-93)		DEPARTMENT OF COMMERCE PATENT AND TRADEMARK OFFICE	ATTORNEY DOCKET NO. 00177/530985MS
TRANSMITTAL LETTER TO THE UNITED STATES DESIGNATED/ELECTED OFFICE (DO/EO/US) CONCERNING A FILING UNDER 35 USC 371			U.S. APPLICATION NO. (if known, see 37 CFR 1.52) NEW 09/508435
International Application No. PCT/JP98/04063	International Filing Date September 10, 1998	Priority Date Claimed September 11, 1997	
Title of Invention IMMUNOASSAY FOR BNP			
Applicant(s) For DO/EO/US Hidehisa ASADA, Hiroyuki SHIMIZU and Kazuaki ENDOU			
Applicant herewith submits to the United States Designated/Elected Office (DO/EO/US) the following items and other information:			
1. <input checked="" type="checkbox"/> This is a FIRST submission of items concerning a filing under 35 USC 371.			
2. <input type="checkbox"/> This is a SECOND or SUBSEQUENT submission of items concerning a filing under 35 USC 371.			
3. <input type="checkbox"/> This is an express request to begin national examination procedures (35 USC 371(f)) at any time rather than delay examination until the expiration of the applicable time limit set in 35 USC 371(b) and PCT Articles 22 and 39(1).			
4. <input checked="" type="checkbox"/> A proper Demand for International Preliminary Examination was made by the 19th month from the earliest claimed priority date.			
5. <input checked="" type="checkbox"/> A copy of the International Application as filed (35 USC 371(c)(2))			
a. <input type="checkbox"/> is transmitted herewith (required only if not transmitted by the International Bureau).			
b. <input checked="" type="checkbox"/> has been transmitted by the International Bureau.			
c. <input type="checkbox"/> is not required, as the application was filed in the United States Receiving Office (RO/US).			
6. <input checked="" type="checkbox"/> A translation of the International Application into English (35 USC 371(c)(2)). ATTACHMENT A			
7. <input type="checkbox"/> Amendments to the claims of the International Application under PCT Article 19 (35 USC 371(c)(3)).			
a. <input type="checkbox"/> are transmitted herewith (required only if not transmitted by the International Bureau).			
b. <input type="checkbox"/> have been transmitted by the International Bureau.			
c. <input type="checkbox"/> have not been made; however, the time limit for making such amendments has NOT expired.			
d. <input type="checkbox"/> have not been made and will not be made.			
8. <input type="checkbox"/> A translation of the amendments to the claims under PCT Article 19 (35 USC 371(c)(3)).			
9. <input checked="" type="checkbox"/> An executed oath or declaration of the inventor(s) (35 USC 371(c)(4)). ATTACHMENT B			
10. <input type="checkbox"/> A translation of the annexes to the International Preliminary Examination Report under PCT Article 36 (35 USC 371(c)(5)).			
Items 11. to 16. below concern other document(s) or information included:			
11. <input checked="" type="checkbox"/> An Information Disclosure Statement under 37 CFR 1.97 and 1.98. ATTACHMENT C			
12. <input checked="" type="checkbox"/> An assignment document for recording. A separate cover sheet in compliance with 37 CFR 3.28 and 3.31 is included. ATTACHMENT D			
13. <input checked="" type="checkbox"/> A FIRST preliminary amendment. ATTACHMENT E			
<input type="checkbox"/> A SECOND or SUBSEQUENT preliminary amendment.			
14. <input type="checkbox"/> A substitute specification.			
15. <input type="checkbox"/> A change of power of attorney and/or address letter.			
16. <input type="checkbox"/> Other items or information:			

THE COMMISSIONER IS AUTHORIZED
TO CURE ANY DEFICIENCY IN THE
FEE FOR THIS PAPER TO DEPOSIT
ACCOUNT NO., 23-0975-

U.S. APPLICATION NO. 09/508435 NEW		INTERNATIONAL APPLICATION NO. PCT/JP98/04063		ATTORNEY DOCKET NO. 00177/530985MS	
17. [X] The following fees are submitted				CALCULATIONS	PTO USE ONLY
BASIC NATIONAL FEE (37 CFR 1.492(a)(1)-(5)):					
[X] Search Report has been prepared by the EPO or JPO \$840.00					
[] International preliminary examination fee paid to USPTO (37 CFR 1.482) \$670.00					
[] No international preliminary examination fee paid to USPTO (37 CFR 1.482) but international search fee paid to USPTO (37 CFR 1.445(a)(2)) \$690.00					
[] Neither international preliminary examination fee (37 CFR 1.482) nor international search fee (37 CFR 1.445(a)(2)) paid to USPTO \$970.00					
[] International preliminary examination fee paid to USPTO (37 CFR 1.482) and all claims satisfied provisions of PCT Article 33(2)-33(4) \$ 96.00					
ENTER APPROPRIATE BASIC FEE AMOUNT =				\$840.00	
Surcharge of \$130.00 for furnishing the oath or declaration later than [] 20 [] 30 months from the earliest claimed priority date (37 CFR 1.492(e)).				\$	
Claims	Number Filed	Number Extra	Rate		
Total Claims	- 20 =		X \$18.00	\$	
Independent Claims	- 3 =		X \$78.00	\$	
Multiple dependent claim(s) (if applicable)			+ \$260.00	\$	
TOTAL OF ABOVE CALCULATIONS =				\$840.00	
Reduction by 1/2 for filing by small entity, if applicable. Verified Small Entity Statement must also be filed. (Note 37 CFR 1.9, 1.27, 1.28)				\$	
SUBTOTAL =				\$840.00	
Processing fee of \$130.00 for furnishing the English translation later than [] 20 [] 30 months from the earliest claimed priority date (37 CFR 1.492(f)).				+	\$
TOTAL NATIONAL FEE =				\$840.00	
Fee for recording the enclosed assignment (37 CFR 1.21(h)). The assignment must be accompanied by an appropriate cover sheet (37 CFR 3.28, 3.31) (\$40 per property).				+	\$40.00
TOTAL FEES ENCLOSED =				\$880.00	
				Amount to be refunded:	\$
				charged:	\$

- a. ☒ A check in the amount of \$880.00 to cover the above fees is enclosed.
- b. ☐ Please charge my Deposit Account No. 23-0975 in the amount of \$_____ to cover the above fees. A duplicate copy of this sheet is enclosed.
- c. ☒ The Commissioner is hereby authorized to charge any additional fees which may be required, or credit any overpayment to Deposit Account No. 23-0975. A duplicate copy of this sheet is enclosed.

NOTE: Where an appropriate time limit under 37 CFR 1.494 or 1.495 has not been met, a petition to revive (37 CFR 1.137(a) or (b)) must be filed and granted to restore the application to pending status.

SEND ALL CORRESPONDENCE TO:

WENDEROTH, LIND & PONACK
2033 K St., N.W., Suite 800
Washington, D.C. 20006

Telephone: (202) 721-8200
Facsimile: (202) 721-8250

By Warren M. Cheek, Jr.
Warren M. Cheek, Jr.
Registration No. 33,367

March 13, 2000
WMC/dlk

Check No. 37182
2000-0259*

5(PRTS

514 P

09/508435
EGIPTO 13 MAR 2000

DESCRIPTION

IMMUNOASSAY FOR BNP

5 TECHNICAL FIELD

The present invention relates to an immunoassay for the brain natriuretic peptide (BNP) which is a member of natriuretic peptide family, more specifically, it relates to an immunoassay for γ -BNP and derivatives thereof.

10

BACKGROUND ART

Natriuretic peptide family includes three members, i.e., atrial natriuretic peptide (ANP), brain natriuretic peptide (BNP) and type C natriuretic peptide (CNP). Among them, ANP and BNP are cardiac hormones which are mainly biosynthesized in and secreted from the heart. ANP and BNP are similar in structure. ANP is a peptide of 28 amino acids with a ring (circular) structure formed by a disulfide bond between the 7th and the 23rd cysteine residues, while BNP is a peptide of 32 amino acids with a ring structure formed by a disulfide bond between the 10th and the 26th cysteine residues. These mature peptides of 28 and 32 amino acids have been considered to be produced from respective precursor when a leader sequence is cleaved off intracellularly or at the time of secretion. That is, there has been reported that human BNP is first synthesized as a preprohormone (hereinafter, referred to as prepro-BNP) in myocardial cells, which is split before or at the time of secretion between Ser²⁶-His²⁷ to give pro-BNP (hereinafter, referred to as γ -BNP), and which is further split between Arg¹⁰²-Ser¹⁰³ to give BNP-32 (hereinafter, referred to as α -BNP) and BNP(1-76), and that the former exhibits the activity. It

has been considered that at least the ring structure must be remained for the expression of activity.

The secretion of cardiac hormones being stimulated by various heart diseases, it well reflects the change in the cardiac functions. The secretion of ANP is accelerated mainly when the atrium undergoes a load, while the biosynthesis and secretion of BNP are stimulated when the ventricle undergoes a load. Accordingly, both ANP and BNP are useful as indicators in the diagnosis of heart disease. As the progress of investigation in the *in vivo* role of respective hormone, the advantageous features of BNP as an indicator for diagnosing heart disease have become clear. For example, the blood concentration of BNP is only 1/6 of ANP in a normal subject but it becomes higher than ANP in patients of heart failure or the like; the blood concentration of BNP increases in the case of heart failure like ANP, and the plasma concentration of BNP often exceeds that of ANP reflecting more accurately the severity of heart dysfunction; the plasma concentration of both ANP and BNP elevates in peripheral blood and elevation rate is higher in BNP. Moreover, BNP level in patients of heart failure sometimes increases to several tens times to several hundreds times of that of healthy normal subjects, and the change of BNP in the cases of heart failure is so marked that no other hormones are incomparable therewith. For these reasons, the usefulness of BNP measurement has been suggested (Y. Saito et al., Mebio, 12(5), 28, 1995).

Under the conditions, an immunoassay which utilizes anti-BNP antibody and is applicable to the diagnosis of cardiac insufficiency has been proposed. Japanese Patent Publication (KOHYO) 7-507210 describes a method of measuring γ -BNP (1-76) produced by biodegradation by protease or the like. However, this

method is directed to γ -BNP (1-76) which lacks the portion(s) essential for the expression of activity such as ring structure and, therefore, cannot determine the hormone activity directly.

5 An assay kit for the measurement of α -BNP having natriuretic activity has been marketed ("BNP-32", Peninsula). With this kit, degradation products of α -BNP in blood including fragments lacking activity due to the deletion of C-terminal region can also be measured. Taking the low blood concentration of BNP into consideration, the measurements involving the degradation products
10 cannot be disregard. Accordingly, the said method connotes disadvantages to be an assay for BNP in the establishment of an accurate diagnosis of heart failure.

As a kit for the measurement of BNP free from the disadvantages above has been marketed ("SHIONORIA BNP",
15 Shionogi), which characteristically uses an antibody recognizing the structure essential for the expression of activity. However, this method would be affected significantly by the process for collecting and storing blood sample, because α -BNP is extremely instable in collected blood. It is, therefore, suggested that the sample should be
20 specifically treated by, for instance, adding an agent for inhibiting degradation into a blood collecting tube or maintaining the sample at low temperature so as to obtain reliable data. Such procedures may hamper the extensive clinical application of the said BNP assay kit.

25 DISCLOSURE OF INVENTION

The present inventors have conducted research intensively for the purpose of establishing an accurate method of diagnosing cardiac diseases involving BNP and found that BNP exists in blood in the form of γ -BNP or its degradation product which at least

comprises structurally the α -BNP moiety (hereinafter, they are referred to as " γ -BNP derivative"), and not in the form of α -BNP which has so far been considered to be dominant. The inventors have also found that γ -BNP is more stable than α -BNP in blood, that is, one role of the N-terminal structure of γ -BNP, among many, would be the stabilization of BNP. The above indicates that an organism biosynthesizes at least 2 kinds of BNP molecule which share the BNP activity but differ in half-life. These findings led the present inventors to have a view that it is indispensable to establish a method specific for not only α -BNP but also γ -BNP to accomplish an accurate diagnosis of cardiac diseases.

The present invention provides an immunoassay specific for mammalian γ -BNP derivatives, characterized in that it uses the first antibody which is reactive with mammalian α -BNP and the second antibody which is reactive with mammalian prepro-BNP or γ -BNP derivatives and not reactive with α -BNP.

As used herein, the term "mammalian α -BNP" refers to a peptide of low molecular weight having natriuretic activity which is derived from mammalian prepro-BNP or γ -BNP through the removal of N-terminal region as a result of processing at the carboxy terminus of processing signal sequence. In case of human BNP, α -BNP is a peptide consisting of C-terminal 32 amino acids (Nos. 103-134) of the amino acid sequence of SEQ ID NO: 1 and having a ring structure. The carboxy terminus of processing signal sequence on the prepro-BNP molecule varies slightly depending on the species. For example, it is No. 102 Arg in case of human BNP while it is No. 100 amino acid in case of porcine or canine BNP.

As used herein, the term "mammalian γ -BNP" refers to a pro-BNP comprising a partial peptide of 32 amino acids corresponding

to α -BNP at the carboxy terminal region. In case of human γ -BNP, it is pro-BNP of 108 amino acids from No. 27 His to No. 134 His of the amino acid sequence of SEQ ID NO: 1. The term "prepro-BNP" refers to a peptide of 134 amino acids from No. 1 Met to No. 134 His of the amino acid sequence of SEQ ID NO: 1 in case of human.

As used herein, the term "mammalian γ -BNP derivative" refers to a peptide fragment derived from mammalian prepro-BNP or γ -BNP through mainly the *in vivo* protease reaction, which fragment includes or is larger than α -BNP. Although γ -BNP derivative would comprise a molecule of the same or smaller size compared to γ -BNP in general, it may comprise a molecule larger than γ -BNP.

Otherwise specifically mentioned, as used herein, the term " γ -BNP derivative" includes γ -BNP itself.

The term "stable", when used herein in connection with BNP, means that a BNP molecule maintains the C-terminal ring structure including C-terminus of BNP and the natriuretic activity after undergoing the degradation by protease, and that the said activity is not significantly decreased even 24 hours from the collection of blood samples. In light of this definition, the γ -BNP derivative as the target substance (analyte) of the present immunoassay is stable.

On the other hand, the term "unstable" means that a BNP sample undergoes degeneration by protease at the C-terminal region and that the natriuretic activity is significantly decreases 24 hours from the collection of blood samples. In light of this definition, α -BNP is unstable.

BRIEF DESCRIPTION OF DRAWINGS

Fig. 1 is a chromatogram obtained in an α -BNP assay system wherein gel filtration HPLC was conducted using Superdex 75 in a

plasma sample. In Fig. 1, A indicates the position of elution of α -BNP.

Fig. 2 is a chromatogram obtained in an α -BNP assay system wherein gel filtration HPLC was conducted using Superdex 75 in a plasma sample different from that shown in Fig. 1. In Fig. 2, A indicates the position of elution of α -BNP.

Fig. 3 is a chromatogram obtained in an immunoassay specific for γ -BNP wherein gel filtration HPLC was conducted using Superdex 75 in a plasma sample same as those shown in Fig. 2. In Fig. 3, A indicates the position of elution of α -BNP.

Fig. 4 is a graph showing the relationships between the storing time and BNP immunoreactivity of γ -BNP kept in human plasma at 25°C.

Fig. 5 is a graph showing the relationships between the storing time and BNP immunoreactivity of α -BNP kept in human plasma at 4°C.

BEST MODE FOR CARRING OUT THE PRESENT INVENTION

In one embodiment of the present invention, it is related to a method which uses two antibodies, wherein the first antibody is reactive with mammalian α -BNP and the second antibody is reactive with prepro-BNP or γ -BNP derivatives and is not reactive with α -BNP.

Antibodies used in the present method can be monoclonal or polyclonal antibodies. The first antibody can be prepared according to a method known in the art using as an antigen human α -BNP which is commercially available or chemically synthesized, or a partial peptide thereof. Alternatively, a monoclonal antibody appended to a commercially available α -BNP assay system (kit) for measuring α -

BNP ("SHIONORIA", Shionogi) is also available, which is reactive with the C-terminal region of α -BNP.

As the second antibody, any antibody can be used subject that it meets the conditions above. Preferred examples of such
5 antibody include those specific for the amino acid sequence shown by the amino acid Nos. 27-102 of SEQ ID NO: 1 or metabolites thereof.

The γ -BNP derivatives as an analyte to be measured by the present method preferably include at least the partial amino acid sequence shown by the amino acid Nos. 27-134 of SEQ ID NO: 1, in case of
10 human BNP. Accordingly, in a preferred embodiment of the present invention, a special attention is preferably paid in the selection of an antigen to obtain an antibody capable of recognizing the amino acid sequence shown by amino acid Nos. 27-102. The preparation of such an antibody can be carried out by any one of methods known in the art.
15 Theoretically, γ -BNP molecule can be cleaved by protease at sites corresponding to No. 47 (Arg), No. 53 (Lys) and No. 72 (Arg) in the amino acid sequence of SEQ ID NO: 1, and, therefore, an antibody recognizing an amino acid sequence shown by amino acid Nos. 73-102 of SEQ ID NO: 1 can be used as the second antibody.

20 The assay of the present invention can be either a competitive- or sandwich-assay and an antibody to be used may be a monoclonal- or polyclonal-antibody.

At least one of the first and the second antibodies may be labeled detectably or immobilized on a solid support.

25 The method for labeling or immobilizing an antibody is known to one ordinary skilled in the art. Examples of label include without limitation radioactive isotopes, enzymes, fluorescent substances, luminescent substances, and particles. The labeling of an antibody can be carried out according to a method known to one ordinary

skilled in the art, for example, that described by Kono et al. (Kaku-Igaku Gijutu, 13(1), 2, (1993)).

The present invention further provides a kit for immunoassay specific to mammalian γ -BNP derivatives, characterized in that it
5 comprises two antibodies wherein the first antibody is reactive with mammalian α -BNP and the second antibody is reactive with mammalian prepro-BNP or γ -BNP derivatives and is not reactive with α -BNP.

The kit of the present invention can be for a competitive- or
10 sandwich-assay and an antibody to be used may be a monoclonal- or polyclonal-antibody.

At least one of the first and the second antibodies may be labeled detectably or immobilized on a solid support. The kit of the present invention may further contain a means for detecting the label.
15 Examples of label include without limitation radioactive isotopes, enzymes, fluorescent substances, luminescent substances, or particles.

The following examples and test examples are provided to further illustrate the present invention, without limiting the scope
20 thereof.

Example 1

Measurement of γ -BNP Derivatives by Sandwich IRMA

Throughout the following Examples, the ordinary reagents used are of special grade supplied by Wako Pure Chemicals Industries, Ltd. or Nacalai Tesque, Inc. The bovine serum albumin (BSA) was
25 purchased from Sigma.

(1) Preparation of Plasma Sample

1) Venous blood was collected from patients of cardiac disease or healthy volunteers and placed in blood-collecting tubes

containing EDTA and aprotinin (500 KIU/l, Sigma) derived from bovine lung. The tubes were centrifuged ($\times 2000$ g at 4°C) for 5 minutes with H-107RGA (Kokusan) to separate blood cells. The resultant plasma samples were freezed and stored at -80°C until use.

5 2) The plasma samples prepared in 1) above from patients of cardiac disease or healthy volunteers were fractionated by gel filtration HPLC system LC10A (Shimadzu) equipped with Superdex 75 10/30 column (Pharmacia). After equilibrating the column with 0.1 M phosphate buffer (pH 7.5, 0.3M NaCl, 5 mM EDTA) at flow rate
10 of 1 ml/min, 1 ml of plasma sample was injected and 1 ml each of effluent eluted from the column was collected. Each fraction was subjected to the measurement by assay systems for measuring α -BNP or γ -BNP as described in (2)-2) and (2)-3) below, respectively.

(2) Construction of Assay System For Measuring α -BNP- or γ -
15 BNP Derivative

1) In the assay system, the following peptides, antibodies and kits were used.

- Human α -BNP (Peptide Institute)
- Antibody against the amino terminal region of γ -hBNP
20 (amino acid Nos. 27-64 of SEQ ID NO: 1) (Peptide Institute)
- Monoclonal antibody against the carboxy terminal structure of α -BNP (BC203). BC203 is an immobilized antibody appended to SIONORIA BNP kit (Shionogi), wherein a monoclonal antibody directed to the carboxy terminal structure of α -BNP is
25 immobilized on beads.

- Monoclonal antibody against the ring structure of α -BNP (KYBNPII). KYBNPII is a monoclonal antibody appended to the SIONORIA BNP kit (Shionogi), which is directed to the ring structure (112-128) of α -BNP, and is labeled with ^{125}I .

2) Measurement of Plasma Fraction by Assay System for α -BNP

The measurement of α -BNP was carried out by commercially available "SHIONORIA BNP kit" (Shionogi). The assay is based on sandwich IRMA (Immunoradiometric Assay) which uses a monoclonal antibody KYBNPII specific for the ring-structure of α -BNP and another monoclonal antibody BC203 specific for the carboxy terminal structure of α -BNP. The assay was carried out in accordance with the supplier's instructions.

That is, 100 μ l each of samples to be assayed or standard solutions (0, 4, 10, 150, 600 or 2000 pg/ml of α -BNP solution) were dispensed into a polystyrene test tube. To the test tube was added 200 μ l of iodine-labeled anti-BNP antibody (125 I) solution, followed by one polystyrene bead on which anti-BC203 antibody has been immobilized. The mixture was stirred and allowed to react by leaving stand for 18 hours at 4 °C. After washing twice with 2 ml of washing solution, radioactivity was measured on γ -counter ARC-600 (Aloka). The results are shown in Figs. 1 and 2.

3) Measurement of Plasma Fraction by Assay System For γ -BNP Derivative

An antibody against amino terminal portion (Nos. 27-64) of γ -hBNP was first labeled with 125 I.

IgG was purified from anti-serum (Peptide Institute) raised against amino terminal portion (amino acid Nos. 27-64 of SEQ ID NO: 1) of γ -hBNP using MASPII kit (Bio-Rad) and displaced with 0.5 M phosphate buffer (pH 7.5) using Centricon 30 (Amicon). The labeling of antibody was carried out by the chloramine T method. To a glass tube was dispensed 170 μ l of purified IgG solution (77.6 μ g, IgG), and 10 μ l of Na 125 I solution (34.2 MBq, Amersham) was added. After

addition of 0.1 % chloramine T solution (20 μ l), the mixture was vigorously stirred at room temperature for 30 seconds. The reaction was quenched by adding 20 μ l of 0.25 % sodium pyrosulfite solution and 20 μ l of 5% aqueous potassium iodide solution. When the
 5 reaction mixture was treated with Ampure SA column (Amersham) to remove unreacted ^{125}I and to desalt, solution containing ^{125}I -labeled antibody was obtained.

The sandwich IRMA was then carried out in plasma fractions by using the resultant antibody and polystyrene beads on which an
 10 antibody recognizing the carboxy terminal structure of α -BNP (BC203).

To a polystyrene tube was placed 100 μ l each of samples to be assayed, followed by 200 μ l of 0.1 M phosphate buffer (pH 7.5, 0.3M, 5 mM EDTA, 0.2% BSA and 500 KIU/l bovine lung aprotinin
 15 (Sigma)) and one polystyrene bead on which BC203 antibody has been immobilized. The mixture was stirred and allowed to react by leaving stand for 18 hours at 4 °C. After washing twice with 2 ml of washing solution, 300 μ l of ^{125}I -labeled antibody solution was added. The mixture was stirred and allowed to react by leaving stand for 18 hours
 20 at 4 °C. After washing twice with 2 ml of washing solution, radioactivity was measured on γ -counter ARC-600 (Aloka) The results are shown in Fig. 3.

(3) Results

Figs. 1, 2 and 3 show the chromatograms of gel filtration
 25 HPLC of plasma samples obtained from patients, wherein A is the position of elution of α -BNP.

Fig. 1 shows the result of the measurement conducted by the α -BNP assay kit described in (2)-2) above. In the Fig. 1, the vertical axis represents the concentration of BNP-like substances in each

fraction and the horizontal axis the volume of effluent eluted from the column as measured by SHINORIA BNP kit. The solid triangle, open square, and open rhombus respectively represent the measurements in different plasma samples.

5 Fig.2 shows the result of the measurement conducted by the α -BNP assay kit described in (2)-2) above in samples different from those shown in Fig. 1. In the Fig. 2, the vertical axis represents the concentration of BNP-like substances in each fraction and the horizontal axis the volume of effluent eluted from the column as measured by SHINORIA BNP kit. The solid triangle and solid square respectively represent the measurements in different plasma samples.

From Figs. 1 and 2, it is revealed that there exist substances of molecular weight larger than α -BNP and having BNP-like immunoreactivity in the plasma of patients of cardiac disease, and that they are the major substances having BNP immunoreactivity.

Fig.3 shows the results of the measurement conducted by γ -BNP assay kit described in (2)-3) above in the same samples those shown in Fig. 2. In Fig. 3, the vertical axis represents the radioactivity measured by the γ -BNP immunoassay system and the horizontal axis the volume of effluent eluted from the column. The solid circle represents the measurements of α -BNP, which obtained after fractionating human α -BNP solution by HPLC in a similar manner as described in the case of plasma.

From Fig. 3, it is revealed that the immunoassay specific for γ -BNP derivative of the present invention can detect the major substances with BNP immunoreactivity, but cannot α -BNP at all.

The results above indicate that the immunoassay for γ -BNP of the present invention is insensitive to α -BNP but specific to γ -BNP derivatives. Further, it has also been revealed that γ -BNP is

the major substance having BNP immunoreactivity.

Test Example 1

Stability of γ -BNP Derivatives and α -BNP in Plasma

Fractions suspected to contain γ -BNP derivative were collected from those obtained by treating plasma samples collected from patients of cardiac disease by gel filtration HPLC. Venous blood was collected from healthy volunteers using blood-collecting tubes containing EDTA in the absence of bovine lung aprotinin. Plasma samples (the minimum detection limit of α -BNP < 4 pg/ml) were prepared in a manner similar to that described in (1)-1) above. The plasma sample was allowed to stand for 0, 2, 6, 24 hours at room temperature (25°C) after addition of the fraction. The stability of BNP derivative was evaluated by determining the BNP immunoreactivity in the plasma sample by means of SHIONORIA BNP kit for assaying α -BNP.

Separately, the stability of α -BNP was evaluated using a plasma sample prepared by adding chemically synthesized α -BNP to plasma collected from healthy volunteers and standing for 0, 2, 6 and 24 hours at 4°C in the absence of bovine lung aprotinin as described above. The BNP immunoreactivity in the plasma sample was determined by SHIONORIA BNP kit in the same manner as above.

The stability of γ -BNP derivatives and α -BNP in plasma samples are shown in Figs. 4 and 5, respectively.

From Fig. 4, it is revealed that γ -BNP derivatives do not lose significantly the immunoreactivity compared with the initial activity even after 24-hour-standing at 25°C. From Fig. 5, by contrast, it is revealed that α -BNP loses the immunoreactivity to about 40% based on the initial activity after 24-hour-standing at 4°C.

The above results demonstrate that α -BNP is far less stable

compared with γ -BNP derivative in blood and that the latter is much more suited in the diagnosis of cardiac diseases than the former.

INDUSTRIAL APPLICABILITY

5 As mentioned above, the BNP level in patients of heart failure sometimes increases to several tens times to several hundreds times of that of healthy normal subjects, and the change of BNP in the cases of heart failure is so marked that no other hormones are incomparable therewith. For this reason, the usefulness of BNP measurement has
10 been proposed.

The immunoassay of the present invention allows to determine specifically γ -BNP derivatives without measuring α -BNP.

Accordingly, the present immunoassay can be a clinically significant means for diagnosis and prognostic monitoring of heart failure, which
15 leads to conclusion/judgment somehow different from those resulted from conventional BNP assay.

Further, it is herein disclosed for the first time that γ -BNP, which is a target substance to be assayed by the present method, is stable in blood. Therefore, immunoassay of the present invention
20 provides stable and reliable clinical data without being affected by the process of collecting or storing samples, or the time from the collection until measurement. Further, the immunoassay of the present invention does not require any special pretreatments of blood sample and therefore gives clinical data conveniently, thereby
25 contributing to the establishment of highly accurate diagnosis of cardiac diseases.

CLAIMS

Sub
B2

1. An immunoassay specific for mammalian γ -BNP derivatives, characterized in that it uses the first antibody reactive with mammalian α -BNP and the second antibody reactive with mammalian prepro-BNP or γ -BNP derivatives and not α -BNP.

2. The immunoassay of claim 1, wherein the γ -BNP derivatives comprise the amino acid sequence shown by the amino acid Nos. 27-102 of SEQ ID NO: 1.

3. The immunoassay of claim 1, wherein the second antibody is specific for the amino acid sequence shown by the amino acid Nos. 27-102 of SEQ ID NO: 1.

4. The immunoassay of ^{Claim 1} ~~any one of claims 1 to 3~~, wherein at least one of the first and the second antibodies is detectably labeled or immobilized.

5. The immunoassay of ^{Claim 1} ~~any one of claims 1 to 3~~, wherein the detectable label is a radioactive isotope, an enzyme, a fluorescent substance, a luminescent substance, or a particle.

6. A kit for immunoassay specific for mammalian γ -BNP derivatives, characterized in that it comprises the first antibody reactive with mammalian α -BNP and the second antibody reactive with mammalian prepro-BNP or γ -BNP derivatives and not α -BNP.

7. The kit of claim 6, wherein at least one of the first and the second antibodies is detectably labeled or immobilized.

8. The kit of claim 7 which further comprises a means for detecting the label.


add
B3 →

ABSTRACT

An immunoassay specific for mammalian γ -BNP derivatives, which uses the first antibody reactive with mammalian α -BNP and
5 the second antibody reactive with mammalian prepro-BNP or γ -BNP derivatives and not α -BNP, and at least one of the first and the second antibodies is optionally labeled detectably labeled or immobilized.

1/5


Fig. 1



2000 21 OCTOBER 2000


2/5

Fig. 2



3/5


Fig. 3



APPROVAL	FIG.
OR	LESS SUBCLASS
RAFTSMAN	


4/5

Fig. 4



5/5

Fig. 5



DECLARATION AND POWER OF ATTORNEY FOR U.S. PATENT APPLICATION

() Original () Supplemental () Substitute (x) PCT () Design

As a below named inventor, I hereby declare that: my residence, post office address and citizenship are as stated below next to my name; that I verily believe that I am the original, first and sole inventor (if only one name is listed below) or an original, first and joint inventor (if plural inventors are named below) of the subject matter which is claimed and for which a patent is sought on the invention entitled:

Title: "IMMUNOASSAY FOR BNP"

of which is described and claimed in:

- () the attached specification, or
 () the specification in the application Serial No. _____ filed _____;
 and with amendments through _____ (if applicable), or
 (x) the specification in International Application No. PCT/ JP98/04063, filed 10.09.98, and as amended
 on _____ (if applicable).

I hereby state that I have reviewed and understand the content of the above-identified specification, including the claims, as amended by any amendment(s) referred to above.

I acknowledge my duty to disclose to the Patent and Trademark Office all information known to me to be material to patentability as defined in Title 37, Code of Federal Regulations, §1.56.

I hereby claim priority benefits under Title 35, United States Code, §119 (and §172 if this application is for a Design) of any application(s) for patent or inventor's certificate listed below and have also identified below any application for patent or inventor's certificate having a filing date before that of the application on which priority is claimed:

COUNTRY	APPLICATION NO.	DATE OF FILING	PRIORITY CLAIMED
Japan	246684/1997	11.09.97	Yes

I hereby claim the benefit under Title 35, United States Code, §120 of any United States application(s) listed below and, insofar as the subject matter of each of the claims of this application is not disclosed in the prior United States application in the manner provided by the first paragraph of Title 35, United States Code, §112, I acknowledge the duty to disclose information material to patentability as defined in Title 37, Code of Federal Regulations, §1.56 which occurred between the filing date of the prior application and the national or PCT international filing date of this application.

APPLICATION SERIAL NO.	U.S. FILING DATE	STATUS: PATENTED, PENDING, ABANDONED

7 And I hereby appoint John T. Miller, Reg. No. 21,120; Michael R. Davis, Reg. No. 25,134; Matthew M. Jacob, Reg. No. 25,154; Jeffrey Noltan, Reg. No. 25,408; Warren M. Cheek, Jr., Reg. No. 33,367; Nils E. Pedersen, Reg. No. 33,145 and Charles R. Watts, Reg. No. 33,142, who together constitute the firm of WENDEROTH, LIND & PONACK, L.L.P., attorneys to prosecute this application and to transact all business in the U.S. Patent and Trademark Office connected therewith.

I hereby authorize the U.S. attorneys named herein to accept and follow instructions from AOYAMA & PARTNERS as to any action to be taken in the U.S. Patent and Trademark Office regarding this application without direct communication between the U.S. attorneys and myself. In the event of a change in the persons from whom instructions may be taken, the U.S. attorneys named herein will be so notified by me.

Send Correspondence to

WENDEROTH, LIND & PONACK, L.L.P.
2033 K Street, N.W., Suite 800
Washington, DC 20006

Direct Telephone Calls to:

WENDEROTH, LIND & PONACK, L.L.P.
Area Code (202) 721-8200

Direct Facsimile Messages to:

Area Code (202) 721-8250

100 Full Name of First Inventor	FAMILY NAME <u>ASADA</u>	FIRST GIVEN NAME <u>Hidehisa</u>	SECOND GIVEN NAME
Residence & Citizenship	CITY <u>Takatsuki</u>	STATE OR COUNTRY <u>Osaka</u>	COUNTRY OF CITIZENSHIP <u>Japan JPX</u>
Post Office Address	ADDRESS <u>5-23-5, Nanpeidai,</u>	CITY <u>Takatsuki-shi,</u>	STATE OR COUNTRY <u>OSAKA</u> ZIP CODE <u>569-1042</u>
200 Full Name of Second Inventor	FAMILY NAME <u>SHIMIZU</u>	FIRST GIVEN NAME <u>Hiroyuki</u>	SECOND GIVEN NAME
Residence & Citizenship	CITY <u>Kobe</u>	STATE OR COUNTRY <u>Hyogo</u>	COUNTRY OF CITIZENSHIP <u>Japan JPX</u>
Post Office Address	ADDRESS <u>Santaun-Oyashiki 709,</u> <u>3-1-2, Oyashiki-dori, Nagataku, Kobe-shi,</u>	CITY <u>HYOGO</u>	STATE OR COUNTRY <u>653-0843</u>
300 Full Name of Third Inventor	FAMILY NAME <u>ENDO</u>	FIRST GIVEN NAME <u>Kazuaki</u>	SECOND GIVEN NAME
Residence & Citizenship	CITY <u>Takatsuki</u>	STATE OR COUNTRY <u>Osaka</u>	COUNTRY OF CITIZENSHIP <u>Japan JPX</u>
Post Office Address	ADDRESS <u>3-5-18, Nanpeidai,</u>	CITY <u>Takatsuki-shi,</u>	STATE OR COUNTRY <u>OSAKA</u> ZIP CODE <u>569-1042</u>
Full Name of Fourth Inventor	FAMILY NAME	FIRST GIVEN NAME	SECOND GIVEN NAME
Residence & Citizenship	CITY	STATE OR COUNTRY	COUNTRY OF CITIZENSHIP
Post Office Address	ADDRESS	CITY	STATE OR COUNTRY ZIP CODE

Full Name of Fifth Inventor	FAMILY NAME	FIRST GIVEN NAME	SECOND GIVEN NAME
Residence & Citizenship	CITY	STATE OR COUNTRY	COUNTRY OF CITIZENSHIP
Post Office Address	ADDRESS	CITY	STATE OR COUNTRY ZIP CODE

Full Name of Sixth Inventor	FAMILY NAME	FIRST GIVEN NAME	SECOND GIVEN NAME
Residence & Citizenship	CITY	STATE OR COUNTRY	COUNTRY OF CITIZENSHIP
Post Office Address	ADDRESS	CITY	STATE OR COUNTRY ZIP CODE

Full Name of Seventh Inventor	FAMILY NAME	FIRST GIVEN NAME	SECOND GIVEN NAME
Residence & Citizenship	CITY	STATE OR COUNTRY	COUNTRY OF CITIZENSHIP
Post Office Address	ADDRESS	CITY	STATE OR COUNTRY ZIP CODE

I further declare that all statements made herein of my own knowledge are true, and that all statements on information and belief are believed to be true; and further that these statements were made with the knowledge that willful false statements and the like so made are punishable by fine or imprisonment, or both, under Section 1001 of Title 18 of the United States Code, and that such willful false statements may jeopardize the validity of the application or any patent issuing thereon.

1st Inventor Yoshihisa Asada Date March 1st, 2000
 2nd Inventor Shirohiko Shimizu Date 2 Mar. 2000
 3rd Inventor Kazuaki Endou Date 3 Mar. 2000
 4th Inventor _____ Date _____
 5th Inventor _____ Date _____
 6th Inventor _____ Date _____
 7th Inventor _____ Date _____

The above application may be more particularly identified as follows:

U.S. Application Serial No. _____ Filing Date _____
 Applicant Reference Number _____ Atty Docket No. _____
 Title of Invention _____

SEQUENCE LISTING

<110> Shionogi & Co., Ltd.

<120> Immunoassay for BNP

5 <130> 660984

<140>

<141>

<150> JP 246684/1997

<151> 1997-9-11

10 <160> 2

<117> Word (MS-DOS text)

<210> 1

<211> 436

<212> DNA

15 <213> human

<400> 1

atg gat ccc cag aca gca cct tcc cgc gcg etc ctg etc ctg etc ttc 48

Met Asp Pro Gln Thr Ala Pro Ser Arg Ala Leu Leu Leu Leu Leu Phe

1 5 10 15

20 ttg cat ctg gct ttc ctg gga ggt cgt tcc cac ccg ctg ggc agc ccc 96

Leu His Leu Ala Phe Lue Gly Gly Arg Ser His Pro Leu Gly Ser Pro

20 25 30

ggc tca gcc tcg gac ttg gaa acg tcc ggg tta cag gag cag cgc aac 144

Gly Ser Ala Ser Asp Leu Glu Thr Ser Gly Leu Gln Glu Gln Arg Asn

25 35 40 45

cat ttg cag ggc aaa ctg tcg gag ctg cag gtg gac cag aca tcc ctg 196

His Leu Gln Gly Lys Leu Ser Glu Leu Gln Val Glu Gln Thr Ser Leu

50 55 60

2/3

gag ccc ctc cag gag agc ccc cgt ccc aca ggt gtc tgg aag tcc cgg 244
Glu Pro Leu Gln Glu Ser Pro Arg Pro Thr Gly Val Trp Lys Ser Arg
65 70 75 80

gag gta gcc acc gag ggc atc cgt ggg cac cgc aaa atg gtc ctc tac 292
Glu Val Ala Thr Glu Gly Ile Arg Gly His Arg Lys Met Val Leu Try
85 90 95

acc ctg cgg gca cca cga agc ccc aag atg gtg caa ggg tct ggc tgc 340
Thr Leu Arg Ala Pro Arg Ser Pro Lys Met Val Gln Gly Ser Gly Cys
100 105 110

ttt ggg agg aag atg gac cgg atc agc tcc tcc agt gcc ctg ggc tgc 388
Phe Gly Arg Lys Met Asp Arg Ile Ser Ser Ser Ser Gly Leu Gly Cys
115 120 125

aaa gtg ctg agg cgg cat 436
Lys Val Leu Arg Arg His
130 134

<210> 2

<211> 134

<212> PRT

<213> human

<400> 2

Met Asp Pro Gln Thr Ala Pro Ser Arg Ala Leu Leu Leu Leu Leu Phe
1 5 10 15

Leu His Leu Ala Phe Leu Gly Gly Arg Ser His Pro Leu Gly Ser Pro
20 25 30

Gly Ser Ala Ser Asp Leu Glu Thr Ser Gly Leu Gln Glu Gln Arg Asn
35 40 45

His Leu Gln Gly Lys Leu Ser Glu Leu Gln Val Glu Gln Thr Ser Leu
50 55 60

5441

3/3

Glu Pro Leu Gln Glu Ser Pro Arg Pro Thr Gly Val Trp Lys Ser Arg
65 70 75 80
Glu Val Ala Thr Glu Gly Ile Arg Gly His Arg Lys Met Val Leu Try
85 90 95
Thr Leu Arg Ala Pro Arg Ser Pro Lys Met Val Gln Gly Ser Gly Cys
100 105 110
Phe Gly Arg Lys Met Asp Arg Ile Ser Ser Ser Ser Gly Leu Gly Cys
115 120 125
Lys Val Leu Arg Arg His
130 134

Sub A1

5

10